

Case Report

IMPLANT INSERTION, BONE REGENERATION AND PALATAL CONNECTIVE TISSUE GRAFTING IN MAXILLARY AESTHETIC AREA: A CASE REPORT WITH A 9-YEAR FOLLOW-UP

S. Gandini¹, E.M. Ravidà¹, G. El Haddad² and E. El Haddad²

¹University of Ferrara, Ferrara, Italy;

²Private practice, Turin, Italy

Correspondence to:

Elias El Haddad, DDS

Corso Vittorio Emanuele II, 38

10123, Turin, Italy

Email: elias.h@gmail.com

ABSTRACT

The restoration of tooth loss in the frontal area is challenging. Patients demand immediate restoration for aesthetic reasons. In addition, tooth loss determines alveolar bone remodeling with consequent reduction in height and thickness of the alveolar bone. Several frontal area rehabilitation techniques exist, ranging from prosthetic solutions to immediately loaded implant solutions. In this case, there is often the need to do soft and hard tissue regeneration. These techniques can be performed at the same or different times of implant insertion. A single intervention to insert the implant, regenerate bone, and increase soft tissue in a single surgical session is more complex but highly appreciated by the patient. The present work describes a case report of a patient who benefited from implant placement, guided bone regeneration, and grafting of connective tissue collected from the palate to restore an upper central incisor.

KEYWORDS: *bone, regeneration, maxilla, jaw, crest, alveolus, graft, fixture*

INTRODUCTION

Immediate post-extractive rehabilitation of loss of frontal teeth is challenging due to soft and hard tissue loss and the request for a high aesthetic solution. The therapeutic choice is guided by several factors, such as the possibility of obtaining an implant's primary stability, gingival biotype, presence/absence of vestibular bone thickness/dehiscence, and marginal bone level.

Ridge bone reabsorption after tooth extraction in the frontal area is a common situation that can compromise a proper gingival tissue level around the implant emerging profile. From this point of view, the literature shows how vestibular bone is essential to maintain the vestibular soft tissue stability of the teeth to be treated (1). In addition, it has been identified that different gingival thicknesses are strictly related to the periodontal probing depth, gingival width, and type of teeth (2). Moreover, a correlation between gingival clinical thickness and alveolar bone thickness on the labial side has been demonstrated using CBCT measurements (3); both variables are essential for proper rehabilitation. Some surgical techniques can be used to maintain or restore alveolar bone ridges, such as socket preservation (4), guided bone

Received: 12 September 2023

Accepted: 19 October 2023

Copyright © by LAB srl 2023 ISSN 2975-1276

This publication and/or article is for individual use only and may not be further reproduced without written permission from the copyright holder. Unauthorized reproduction may result in financial and other penalties. Disclosure: All authors report no conflicts of interest relevant to this article.

regeneration, and the tent pole technique (5). Since patients ask for an urgent solution, combining some surgical procedures (i.e., implant insertion, bone regeneration, and soft tissue grafting) in a single surgical time is possible. In these cases, patient selection is paramount since the risk of failure is higher in cases where each surgical procedure is performed at different times.

Palatal tissue graft is the most used technique for soft tissue augmentation. The connective tissue graft procedure requires harvesting soft tissue from the palate (i.e., donor site), freeing the graft from the epithelium, and applying the connective tissue in a recipient site. The choice of the donor site is guided by the quantitative requirements of the site that received the graft (6). Indeed, connective tissue graft increases the width and thickness of keratinized tissue and improves the therapy's aesthetic results (7).

The present work reports a case treated with post-extractive implant insertion, guided bone regeneration using heterologous material, and palatal harvest connective tissue grafting treated by a single surgeon (Dr. Elias El Haddad). Furthermore, pertinent literature is discussed.

CASE REPORT

At the first visit, a 53-year-old man complained about a purulent swelling at the level of the left upper central incisor (Fig. 1).



Fig. 1. *Initial stage, frontal view*

A radiographic examination revealed the development of an apical lesion with probable apicectomy performed in the past (Fig. 2). Dental probing showed no periodontal lesions around the central incisor or adjacent teeth.



Fig. 2. *Radiographic examination with the development of an apical lesion*

The clinician explained different treatment options to the patient. The choice selected by the surgeon and the patient was to extract the tooth and immediately insert an implant 4.3 x 18 mm, replace tapered with a TiUnite surface, Nobel Biocare®, perform a bone regeneration using heterologous material, and graft connective tissue harvest from the palate. After taking a pre-extraction impression, the tooth element was provisionally restored with a temporary resin crown prepared in advance by the dental laboratory.

From a surgical point of view, after performing local anesthesia with adrenaline, the tooth was extracted, and a flap from the upper right central incisor to the upper left cuspid was lifted without a vertical incision. Then, a dehiscence in the vestibular bone plate was visible (Fig. 3).




Fig. 3. *In the first photo, we can see the flap's realization and the vestibular bone's dehiscence in the following photo.*

The alveolus was cleaned by granulation tissue, and the implant was inserted. The implant surface was out of bone in the vestibular area for about 8 mm (Fig. 4).




Fig. 4. *Implant positioning and vestibular dehiscence.*

Notably, the implant's surface was under the vestibular bone profile, a condition that makes bone regeneration ideal. A temporary crown was placed before proceeding with the reconstruction of hard and soft tissues. A cemented temporary crown was chosen to manage the restoration's aesthetics better.

A standard abutment was connected to the implant, leaving a space of 2 mm between it and the lower teeth to have the space for to create a correct occlusal crown. Afterward, our temporary crown was checked for occlusion in a nonfunctional way and finished to prevent over-contours and excess resin. It was polished and then cemented with a layer of temporary cement (Fig. 5).



Fig. 5. *Placement of a cemented temporary crown on the implant.*

Once this phase was completed and the excellent stability of our implant was verified (by tightening and loosening the abutment screw), the surgeon proceeded with the reconstruction of hard and soft tissues. A connective tissue graft was taken from the palate side of the same surgical field. It was unnecessary to reach the molar area to collect the palatine connective tissue, which was preserved in physiological saline solution (Fig. 6).




Fig. 6. *Connective tissue graft taken from the palate*

The vestibular bone defect was filled with a granular graft material of bovine origin, which was then covered with platelet-derived membranes (Fig. 7).




Fig. 7. *Granular graft material of bovine origin and platelet-derived membranes*

Above the bovine-derived granular graft material, the connective graft was positioned and sutured with the primary flap (Fig. 8).




Fig. 8. *Positioning of the connective graft and suture with primary flap.*

An X-ray was performed to check implant positioning and prosthetic rehabilitation, and the patient was discharged with antibiotic therapy (i.e., amoxicillin and clavulanic acid), a chlorhexidine-based mouthwash, and a pain reliever (Fig. 9). One week later, the sutures were removed (Fig. 10).




Fig. 9. Clinical and X-ray control



Fig. 10. Suture removal after one week.

After three months, the provisional restoration was finalized with a ceramic crown (Fig.11). Fig. 12 shows the clinical and radiological checks before and after the treatment.



Fig. 11. Ceramic prosthesis.



Fig. 12. Clinical and radiological check before and after the treatment.

A 9-year follow-up CBCT control was performed to better evaluate the vestibular bone regeneration outcome (Fig. 13).




Fig.13. 9-year follow-up with CBCT investigation. A): Immediately after surgery; B): Nine years after surgery.

DISCUSSION

For a long time, research in the implant field focused on peri-implant bone, as it claimed that quality and quantity of hard tissue were the main criteria for success (8). Therefore, before placing an implant, it was essential to have adequate bone volume, and this could be achieved through various conservation and ridge augmentation procedures (9).

The critical factors for implant success have always been crestal bone loss (10) and primary implant stability (11), while the importance of peri-implant soft tissue has often been overlooked. Recently, scientific evidence has shown that peri-implant soft tissue is critical to maintaining peri-implant health (12). Several publications evaluated the importance of soft tissues at dental implant sites from both a biological (13-15) and esthetic perspective (16, 17).

The anatomy of peri-implant mucosa and gingiva are different according to several factors. First, there is a difference in the connective tissue fibers that do not attach to the implant (18). In addition, the implant site has reduced vascular supply (19), and the junctional epithelium around implants is more permeable (20). Maintaining an adequate quantity and quality of mucosa around peri-implant bone is essential to promote peri-implant health (21).

Connective tissue grafts have become an essential part of periodontal reconstructive surgery, and several techniques are available to harvest a suitable connective tissue graft (22). For immediately loaded implants, strong evidence suggests that their placement and simultaneous bone grafting should be combined with soft tissue grafting to counteract postoperative remodeling processes. Adding a connective tissue graft has been shown to improve aesthetics and reduce the formation of peri-implant soft tissue recessions (23-25).

The reported case with a 9-year follow-up strengthens the importance of connective grafting in conjunction with bone regeneration at the time of implant insertion.

REFERENCES

1. Kois JC, Kan JY. Predictable peri-implant gingival aesthetics: surgical and prosthodontic rationales. *Practical procedures & aesthetic dentistry: PPAD*. 2001;13(9):691-698; quiz 700, 721-722.
2. Muller HP, Eger T. Gingival phenotypes in young male adults. *Journal of Clinical Periodontology*. 1997;24(1):65-71. doi:https://doi.org/10.1111/j.1600-051x.1997.tb01186.x
3. Fu JH, Yeh CY, Chan HL, Tatarakis N, Leong DJM, Wang HL. Tissue Biotype and Its Relation to the Underlying Bone Morphology. *Journal of Periodontology*. 2010;81(4):569-574. doi:https://doi.org/10.1902/jop.2009.090591
4. Fee L. Socket preservation. *British Dental Journal*. 2017;222(8):579-582. doi:https://doi.org/10.1038/sj.bdj.2017.355
5. Daga D, Mehrotra D, Mohammad S, Singh G, Natu SM. Tentpole technique for bone regeneration in vertically deficient alveolar ridges: A review. *Journal of Oral Biology and Craniofacial Research*. 2015;5(2):92-97. doi:https://doi.org/10.1016/j.jobcr.2015.03.001
6. Zucchelli G. Tecniche di prelievo dell'innesto connettivale. In: *Chirurgia Estetica Mucogengivale*. Quintessenza; 2014:424-457.
7. Cairo F, Pagliaro U, Nieri M. Soft tissue management at implant sites. *Journal of Clinical Periodontology*. 2008;35(8S):163-

167. doi:<https://doi.org/10.1111/j.1600-051x.2008.01266.x>
8. Albrektsson T, Buser D, Lars Sennerby. On crestal/marginal bone loss around dental implants. *The International journal of periodontics & restorative dentistry*. 2012;25(4):320-322.
 9. Benic GI, Hämmerle CHF. Horizontal bone augmentation by means of guided bone regeneration. *Periodontology 2000*. 2014;66(1):13-40. doi:<https://doi.org/10.1111/prd.12039>
 10. De Bruyn H, Vandeweghe S, Ruyffelaert C, Cosyn J, Sennerby L. Radiographic evaluation of modern oral implants with emphasis on crestal bone level and relevance to peri-implant health. *Periodontology 2000*. 2013;62(1):256-270. doi:<https://doi.org/10.1111/prd.12004>
 11. Sennerby L, Meredith N. Implant stability measurements using resonance frequency analysis: biological and biomechanical aspects and clinical implications. *Periodontology 2000*. 2008;47(1):51-66. doi:<https://doi.org/10.1111/j.1600-0757.2008.00267.x>
 12. Zigdon H, Machtei EE. The dimensions of keratinized mucosa around implants affect clinical and immunological parameters. *Clinical Oral Implants Research*. 2008;19(4):387-392. doi:<https://doi.org/10.1111/j.1600-0501.2007.01492.x>
 13. Brito C, Tenenbaum HC, Wong BKC, Schmitt C, Nogueira-Filho G. Is keratinized mucosa indispensable to maintain peri-implant health? A systematic review of the literature. *Journal of Biomedical Materials Research Part B: Applied Biomaterials*. 2013;102(3):643-650. doi:<https://doi.org/10.1002/jbm.b.33042>
 14. Lin GH, Chan HL, Wang HL. The Significance of Keratinized Mucosa on Implant Health: A Systematic Review. *Journal of Periodontology*. 2013;84(12):1755-1767. doi:<https://doi.org/10.1902/jop.2013.120688>
 15. Souza AB, Tormena M, Matarazzo F, Araújo MG. The influence of peri-implant keratinized mucosa on brushing discomfort and peri-implant tissue health. *Clinical Oral Implants Research*. 2015;27(6):650-655. doi:<https://doi.org/10.1111/clr.12703>
 16. Brandenburg FD, Sailer I, Fehmer V, Büchi DLE, Hämmerle CHF, Thoma DS. Randomized controlled clinical pilot study of all-ceramic single-tooth implant reconstructions: clinical and microbiological outcomes at one year of loading. *Clinical Oral Implants Research*. 2016;28(4):406-413. doi:<https://doi.org/10.1111/clr.12813>
 17. Ioannidis A, Cathomen E, Jung RE, Fehmer V, Hüsler J, Thoma DS. Discoloration of the mucosa caused by different restorative materials - a spectrophotometric in vitro study. *Clinical Oral Implants Research*. 2016;28(9):1133-1138. doi:<https://doi.org/10.1111/clr.12928>
 18. Berglundh T, Lindhe J, Ericsson I, Marinello CP, Liljenberg B, Thomsen P. The soft tissue barrier at implants and teeth. *Clinical Oral Implants Research*. 1991;2(2):81-90. doi:<https://doi.org/10.1034/j.1600-0501.1991.020206.x>
 19. Berglundh T, Lindhe J, Jonsson K, Ericsson I. The topography of the vascular systems in the periodontal and peri-implant tissues in the dog. *Journal of Clinical Periodontology*. 1994;21(3):189-193. doi:<https://doi.org/10.1111/j.1600-051x.1994.tb00302.x>
 20. Moon IS, Berglundh T, Abrahamsson I, Linder E, Lindhe J. The barrier between the keratinized mucosa and the dental implant. *Journal of Clinical Periodontology*. 1999;26(10):658-663. doi:<https://doi.org/10.1034/j.1600-051x.1999.261005.x>
 21. Nemcovsky CE, Moses O. Rotated palatal flap. A surgical approach to increase keratinized tissue width in maxillary implant uncovering: technique and clinical evaluation. *Int J Periodontics Restorative Dent*. 2002;22(6):607-612.
 22. Harris RJ. A comparison of two techniques for obtaining a connective tissue graft from the palate. *International Journal of Periodontics & Restorative Dentistry*. 1997;17(3):260-271.
 23. Cornelini R, Barone A, Ugo Covani. Connective tissue grafts in postextraction implants with immediate restoration: a prospective controlled clinical study. *Pract Proced Aesthet Dent*. 2008;20(6):337-343.
 24. Cosyn J, Eghbali A, Hermans A, Vervaeke S, De Bruyn H, Cleymaet R. A 5-year prospective study on single immediate implants in the aesthetic zone. *Journal of Clinical Periodontology*. 2016;43(8):702-709. doi:<https://doi.org/10.1111/jcpe.12571>
 25. Zuhr O, Bäumer D, Hürzeler M. The addition of soft tissue replacement grafts in plastic periodontal and implant surgery: critical elements in design and execution. *Journal of Clinical Periodontology*. 2014;41(15S):S123-S142. doi:<https://doi.org/10.1111/jcpe.12185>